


Glutathione oxidized

Cat. No.:	HY-D0844
CAS No.:	27025-41-8
Molecular Formula:	C ₂₀ H ₃₂ N ₆ O ₁₂ S ₂
Molecular Weight:	612.63
Target:	Endogenous Metabolite; Reactive Oxygen Species
Pathway:	Metabolic Enzyme/Protease; Immunology/Inflammation; NF-κB
Storage:	Sealed storage, away from moisture
	Powder -80°C 2 years
	-20°C 1 year
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 250 mg/mL (408.08 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions		1 mg	5 mg	10 mg
		1 mM	1.6323 mL	8.1615 mL	16.3231 mL
5 mM		0.3265 mL	1.6323 mL	3.2646 mL	
	10 mM	0.1632 mL	0.8162 mL	1.6323 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: PBS Solubility: 100 mg/mL (163.23 mM); Clear solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	Glutathione oxidized (GSSG) is produced by the oxidation of glutathione. Detoxification of reactive oxygen species is accompanied by production of glutathione oxidized. Glutathione oxidized can be used for the research of sickle cells and erythrocytes ^[1] .
IC₅₀ & Target	Human Endogenous Metabolite
In Vitro	Glutathione oxidized concentrations can be increased with the addition of (2,3-dimethoxy-1,4-naphthoquinonethe) DMNQ in erythrocytes ^[1] . Glutathione oxidized concentrations of intracellular and extracellular can be increased with the stimulation of H ₂ O ₂ in sickle erythrocytes than in healthy erythrocytes ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Adv Sci (Weinh). 2022 Oct 18;e2203088.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Nur E, et al. Increased efflux of oxidized glutathione (GSSG) causes glutathione depletion and potentially diminishes antioxidant defense in sickle erythrocytes. Biochim Biophys Acta. 2011 Nov;1812(11):1412-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA