


ADH-1 trifluoroacetate

Cat. No.:	HY-13541A
CAS No.:	1135237-88-5
Molecular Formula:	C ₂₄ H ₃₅ F ₃ N ₈ O ₈ S ₂
Molecular Weight:	684.71
Target:	Others
Pathway:	Others
Storage:	Sealed storage, away from moisture
	Powder -80°C 2 years
	-20°C 1 year

* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 43 mg/mL (62.80 mM)
 H₂O : 33.33 mg/mL (48.68 mM; Need ultrasonic)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		1.4605 mL	7.3024 mL	14.6047 mL
	5 mM		0.2921 mL	1.4605 mL	2.9209 mL
	10 mM		0.1460 mL	0.7302 mL	1.4605 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 50 mg/mL (73.02 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (3.65 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (3.65 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (3.65 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

ADH-1 trifluoroacetate is an N-cadherin antagonist, which inhibits N-cadherin mediated cell adhesion.

In Vitro

ADH-1 (0.2 mg/mL) blocks collagen I-mediated changes in pancreatic cancer cells, and is highly effective at preventing cell

motility that is induced by expression of N-cadherin. ADH-1 (0, 0.1, 0.2, 0.5 and 1.0 mg/mL) induces apoptosis in a dose-dependent and N-cadherin-dependent manner^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

ADH-1 (50 mg/kg) significantly prevents tumor growth and metastasis in a mouse model for pancreatic cancer. ADH-1 prevents tumor cell invasion and metastasis in an orthotopic model for pancreatic cancer using N-cadherin overexpressing BxPC-3 cells^[1]. ADH-1, at the dosages evaluated, does not display either antiangiogenic activity in a rat aortic ring assay or antitumor potential in a PC3 subcutaneous xenograft tumor model^[2]. ADH-1 (10 mL/kg, i.p.) augmentation of melanoma tumor growth is overcome through its ability to make regionally infused melphalan more effective. ADH-1 mediated augmentation of melanoma tumor growth is not altered by regionally infused temozolomide. In A375, but not DM443 xenografts, ADH-1 treatment increases phosphorylation of AKT at serine 473. ADH-1 slightly diminishes N-cadherin expression in both xenografts^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Animal Administration ^[1]

Animals are anesthetized, and 40 μ L of a single cell suspension containing 50,000 cells is injected into the pancreas. Mice are randomized into treatment groups 10 days after surgery. For treatment, mice are injected intraperitoneally once per day with ADH-1 at 50 mg/kg in 100 μ L PBS (\times 1 per day, \times 5 per week for 4 weeks). For in vivo bioluminescence, D-Luciferin is administered by intraperitoneal injection. Data are acquired 20 min after injection using the IVIS system. Tumor growth is monitored every 10 days from day 10 to day 50 after surgery. Luciferase activity is quantified using the IVIS system. Two months after surgery, the mice are killed, and the pancreas, liver, lung, and disseminated nodules are harvested, fixed in 10% buffered formalin, and embedded in paraffin. Serial 5- μ m sections are cut, mounted on slides, and stained with H&E using standard procedures. Sections are also stained for TUNEL. Sections are examined using a Zeiss Axioscop 40 microscope equipped with an AxioCam MR digital camera and software.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Oncogene. 2022 Oct 12.
- Cell Death Dis. 2022 Jun 20;13(6):557.
- J Cell Mol Med. 2022 Feb 27.
- Sci Rep. 2019 Feb 6;9(1):1517.
- Biochem Biophys Res Commun. 2019 Oct 5. pii: S0006-291X(19)31851-0.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Shintani Y, et al. ADH-1 suppresses N-cadherin-dependent pancreatic cancer progression. *Int J Cancer*. 2008 Jan 1;122(1):71-7.

[2]. Li H, et al. ADH1, an N-cadherin inhibitor, evaluated in preclinical models of angiogenesis and androgen-independent prostate cancer. *Anticancer Drugs*. 2007 Jun;18(5):563-8.

[3]. Turley RS, et al. Targeting N-cadherin increases vascular permeability and differentially activates AKT in melanoma. *Ann Surg*. 2015 Feb;261(2):368-77

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA