(S)-2-((tert-Butoxycarbonyl)amino)-2-cyclohexylacetic acid

Cat. No.: HY-W002410 CAS No.: 109183-71-3 Molecular Formula: C₁₃H₂₃NO₄ 257.33 Molecular Weight:

Target: **Amino Acid Derivatives**

Pathway: Others

Powder Storage: -20°C 3 years

2 years

In solvent -80°C 6 months

> -20°C 1 month

Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

DMSO: 100 mg/mL (388.61 mM; Need ultrasonic)

| Preparing Stock Solutions | Solvent Mass Concentration | 1 mg | 5 mg | 10 mg |
|------------------------------|-------------------------------|-----------|------------|------------|
| | 1 mM | 3.8861 mL | 19.4303 mL | 38.8606 mL |
| | 5 mM | 0.7772 mL | 3.8861 mL | 7.7721 mL |
| | 10 mM | 0.3886 mL | 1.9430 mL | 3.8861 mL |

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (9.72 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (9.72 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (9.72 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

(S)-2-((tert-Butoxycarbonyl)amino)-2-cyclohexylacetic acid is a <u>Glycine</u> (HY-Y0966) derivative^[1].

In Vitro

Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

| REFERENCES | | | | | | | |
|--|--------------------------|------------------------------|--------------------------------|--------------|--|--|--|
| [1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-1144. | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | Caution: Product has not | been fully validated for med | dical applications. For resear | ch use only. | | | |
| | Tel: 609-228-6898 | Fax: 609-228-5909 | E-mail: tech@MedChemE | xpress.com | | | |
| | Address: 1 D | eer Park Dr, Suite Q, Monmo | uth Junction, NJ 08852, USA | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |

Page 2 of 2 www.MedChemExpress.com