## Fmoc-D-Cys(Trt)-OH

MedChemExpress

HY-W01072	4	
167015-11-4	1	
C <sub>37</sub> H <sub>31</sub> NO <sub>4</sub> S		
585.71		
Amino Acid	Derivativ	es
Others		
Powder	-20°C	3 years
	4°C	2 years
In solvent	-80°C	6 months
	-20°C	1 month
	167015-11-4 C <sub>37</sub> H <sub>31</sub> NO <sub>4</sub> S 585.71 Amino Acid Others Powder	585.71 Amino Acid Derivativ Others Powder -20°C 4°C In solvent -80°C

®

## SOLVENT & SOLUBILITY

	Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg	
		1 mM	1.7073 mL	8.5366 mL	17.0733 mL	
		5 mM	0.3415 mL	1.7073 mL	3.4147 mL	
		10 mM	0.1707 mL	0.8537 mL	1.7073 mL	
	Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (4.27 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (4.27 mM); Clear solution					

<b>BIOLOGICAL ACTIV</b>	ИТҮ
Description	Fmoc-D-Cys(Trt)-OH is a cysteine derivative <sup>[1]</sup> .
In Vitro	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

[1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-807.

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA