(S)-2-((((9H-Fluoren-9-yl)methoxy)carbonyl)amino)-2,3-dimethylbutanoic acid

Cat. No.: HY-W018849

CAS No.: 169566-81-8

Molecular Formula: $C_{21}H_{23}NO_4$

Molecular Weight: 353.42

Target: Amino Acid Derivatives

Pathway: Others

Storage:

Powder	-20°C	3 years
	4°C	2 years
In solvent	-80°C	6 months
	-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (282.95 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
	1 mM	2.8295 mL	14.1475 mL	28.2949 mL
	5 mM	0.5659 mL	2.8295 mL	5.6590 mL
	10 mM	0.2829 mL	1.4147 mL	2.8295 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
   Solubility: ≥ 2.5 mg/mL (7.07 mM); Clear solution
2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
   Solubility: ≥ 2.5 mg/mL (7.07 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 90% corn oil
   Solubility: ≥ 2.5 mg/mL (7.07 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

(S)-2-((((9H-Fluoren-9-yl)methoxy)carbonyl)amino)-2,3-dimethylbutanoic acid is a valine derivative[1].

In Vitro

Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-1144.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA