

GIP, human

Cat. No.:	HY-P0276	
CAS No.:	100040-31-1	
Molecular Formula:	$C_{226}H_{338}N_{60}O_{66}S$	
Molecular Weight:	4983.6	YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGGKNDWKHNITQ
Sequence:	Tyr-Ala-Glu-Gly-Thr-Phe-Ile-Ser-Asp-Tyr-Ser-Ile-Ala-Met-Asp-Lys-Ile-His-Gln-Gln-Asp-Phe-Val-Asn-Trp-Leu-Leu-Ala-Gln-Lys-Gly-Lys-Lys-Asn-Asp-Trp-Lys-His-Asn-Ile-Thr-Gln	
Sequence Shortening:	YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGGKNDWKHNITQ	
Target:	Insulin Receptor	
Pathway:	Protein Tyrosine Kinase/RTK	
Storage:	Sealed storage, away from moisture and light, under nitrogen	
	Powder -80°C 2 years	
	-20°C 1 year	
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)	

SOLVENT & SOLUBILITY

In Vitro

H₂O : 25 mg/mL (5.02 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	0.2007 mL	1.0033 mL	2.0066 mL
	5 mM	0.0401 mL	0.2007 mL	0.4013 mL
	10 mM	---	---	---

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

GIP, human, a peptide hormone consisting of 42 amino acids, is a stimulator of glucose-dependent insulin secretion and a weak inhibitor of gastric acid secretion. GIP, human acts as an incretin hormone released from intestinal K cells in response to nutrient ingestion^{[1][2][3]}.

In Vitro

Gastric Inhibitory Polypeptide (GIP) exerts various peripheral effects on adipose tissue and lipid metabolism, thereby leading to increased lipid deposition in the postprandial state^[1]. GIP, human plays a vital role in lipid metabolism and the development of obesity.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Toxins. 2021, 13(8), 512.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Meier JJ, et al. Gastric inhibitory polypeptide: the neglected incretin revisited. Regul Pept. 2002 Jul 15;107(1-3):1-13.
- [2]. Miyachi A, et al. Quantitative analytical method for determining the levels of gastric inhibitory polypeptides GIP1-42 and GIP3-42 in human plasma using LC-MS/MS/MS. J Proteome Res. 2013;12(6):2690-2699.
- [3]. Gabe MBN, et al. Molecular interactions of full-length and truncated GIP peptides with the GIP receptor - A comprehensive review. Peptides. 2020;125:170224.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA