

Charybdotoxin TFA

Cat. No.:	HY-P0191A
Molecular Formula:	C ₁₇₆ H ₂₇₇ N ₅₇ O ₅₅ S ₇ C ₂ HF ₃ O ₂
Molecular Weight:	4409.91
Sequence:	{Glp}-Phe-Thr-Asn-Val-Ser-Cys-Thr-Thr-Ser-Lys-Glu-Cys-Trp-Ser-Val-Cys-Gln-Arg-Leu-His-Asn-Thr-Ser-Arg-Gly-Lys-Cys-Met-Asn-Lys-Lys-Cys-Arg-Cys-Tyr-Ser (Disulfide bridge: Cys7-Cys28; Cys13-Cys33; Cys17-Cys35)
Sequence Shortening:	{Glp}-FTNVSCTTSKECWSVCQRLHNTSRGKCMNKKCRCYS (Disulfide bridge: Cys7-Cys28; Cys13-Cys33; Cys17-Cys35)
Target:	Potassium Channel
Pathway:	Membrane Transporter/Ion Channel
Storage:	Sealed storage, away from moisture and light Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)

BIOLOGICAL ACTIVITY

Description	Charybdotoxin TFA, a 37-amino acid peptide, is a K ⁺ channel blocker ^[1] .
In Vitro	Charybdotoxin represents a remarkable tool for studying K ⁺ channels ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. M L Garcia , et al. Charybdotoxin and its effects on potassium channels. Am J Physiol. 1995 Jul;269(1 Pt 1):C1-10.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA