

GIP, human TFA

Cat. No.:	HY-P0276A
Molecular Formula:	C ₂₂₈ H ₃₃₉ F ₃ N ₆₀ O ₆₈ S
Molecular Weight:	5097.62
Sequence:	Tyr-Ala-Glu-Gly-Thr-Phe-Ile-Ser-Asp-Tyr-Ser-Ile-Ala-Met-Asp-Lys-Ile-His-Gln-Gln-Asp-Phe-Val-Asn-Trp-Leu-Leu-Ala-Gln-Lys-Gly-Lys-Lys-Asn-Asp-Trp-Lys-His-Asn-Ile-Thr-Gln <small>YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGGKNDWKHNITQ (TFA salt)</small>
Sequence Shortening:	YAEGTFISDYSIAMDKIHQQDFVNWLLAQKGGKNDWKHNITQ
Target:	Insulin Receptor
Pathway:	Protein Tyrosine Kinase/RTK
Storage:	Sealed storage, away from moisture and light, under nitrogen Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)

SOLVENT & SOLUBILITY

In Vitro

H₂O : 10 mg/mL (1.96 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	0.1962 mL	0.9808 mL	1.9617 mL
	5 mM	---	---	---
	10 mM	---	---	---

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

GIP, human TFA, a peptide hormone consisting of 42 amino acids, is a stimulator of glucose-dependent insulin secretion and a weak inhibitor of gastric acid secretion. GIP, human TFA acts as an incretin hormone released from intestinal K cells in response to nutrient ingestion^{[1][2][3]}.

In Vitro

Gastric Inhibitory Polypeptide (GIP) exerts various peripheral effects on adipose tissue and lipid metabolism, thereby leading to increased lipid deposition in the postprandial state^[1]. GIP, human plays a vital role in lipid metabolism and the development of obesity.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Toxins. 2021, 13(8), 512.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Meier JJ, et al. Gastric inhibitory polypeptide: the neglected incretin revisited. Regul Pept. 2002 Jul 15;107(1-3):1-13.
- [2]. Miyachi A, et al. Quantitative analytical method for determining the levels of gastric inhibitory polypeptides GIP1-42 and GIP3-42 in human plasma using LC-MS/MS/MS. J Proteome Res. 2013;12(6):2690-2699.
- [3]. Gabe MBN, et al. Molecular interactions of full-length and truncated GIP peptides with the GIP receptor - A comprehensive review. Peptides. 2020;125:170224.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA