


## Fmoc-Tyr-OH

<b>Cat. No.:</b>	HY-W009003		
<b>CAS No.:</b>	92954-90-0		
<b>Molecular Formula:</b>	C <sub>24</sub> H <sub>21</sub> NO <sub>5</sub>		
<b>Molecular Weight:</b>	403.43		
<b>Target:</b>	Amino Acid Derivatives		
<b>Pathway:</b>	Others		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 66.67 mg/mL (165.26 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	2.4787 mL	12.3937 mL	24.7874 mL
		5 mM	0.4957 mL	2.4787 mL	4.9575 mL
10 mM		0.2479 mL	1.2394 mL	2.4787 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	<p>1. Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: ≥ 2.5 mg/mL (6.20 mM); Clear solution</p> <p>2. Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: ≥ 2.5 mg/mL (6.20 mM); Clear solution</p>				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Fmoc-Tyr-OH is a tyrosine derivative <sup>[1]</sup> .
<b>In Vitro</b>	<p>Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

### REFERENCES

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA