Fmoc-Asp-OtBu

Cat. No.: HY-W010959

CAS No.: 129460-09-9

Molecular Formula: $C_{23}H_{25}NO_6$

Molecular Weight: 411.45

Target: Amino Acid Derivatives

Pathway: Others

Storage:

Powder	-20°C	3 years
	4°C	2 years
In solvent	-80°C	6 months
	-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (243.04 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
	1 mM	2.4304 mL	12.1521 mL	24.3043 mL
	5 mM	0.4861 mL	2.4304 mL	4.8609 mL
	10 mM	0.2430 mL	1.2152 mL	2.4304 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
   Solubility: ≥ 2.5 mg/mL (6.08 mM); Clear solution
2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
   Solubility: ≥ 2.5 mg/mL (6.08 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 90% corn oil
   Solubility: ≥ 2.5 mg/mL (6.08 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Fmoc-Asp-OtBu is an aspartic acid derivative[1].

In Vitro

Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Luckose F, et al. Effects of amino acid derivatives on physical, mental, and physiological activities. Crit Rev Food Sci Nutr. 2015;55(13):1793-1144.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA