


## Fmoc-D-Isoleucine

<b>Cat. No.:</b>	HY-W011135		
<b>CAS No.:</b>	143688-83-9		
<b>Molecular Formula:</b>	C <sub>21</sub> H <sub>23</sub> NO <sub>4</sub>		
<b>Molecular Weight:</b>	353.41		
<b>Target:</b>	Amino Acid Derivatives		
<b>Pathway:</b>	Others		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 100 mg/mL (282.96 mM; Need ultrasonic)				
		<b>Solvent</b>	<b>Mass</b>		
		<b>Concentration</b>	<b>1 mg</b>	<b>5 mg</b>	
	<b>Preparing Stock Solutions</b>		<b>10 mg</b>		
		<b>1 mM</b>	2.8296 mL	14.1479 mL	28.2957 mL
		<b>5 mM</b>	0.5659 mL	2.8296 mL	5.6591 mL
		<b>10 mM</b>	0.2830 mL	1.4148 mL	2.8296 mL
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.07 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.07 mM); Clear solution				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Fmoc-D-Isoleucine is an isoleucine derivative <sup>[1]</sup> .
<b>In Vitro</b>	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA