

## Transferrins

Cat. No.:	HY-P3267
CAS No.:	11096-37-0
Target:	Others
Pathway:	Others
Storage:	Sealed storage, away from moisture and light
	Powder    -80°C    2 years
	-20°C    1 year

\* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)

# Transferrins

### SOLVENT & SOLUBILITY

In Vitro	H <sub>2</sub> O : 100 mg/mL (Need ultrasonic)
----------	--

### BIOLOGICAL ACTIVITY

Description	Transferrin, a blood-plasma glycoprotein, is a target ligand for transferrin receptor. Transferrins can bind to and mediate the transport of iron <sup>[1][2][3]</sup> .
IC <sub>50</sub> & Target	Transferrin receptor <sup>[1]</sup>
In Vitro	<p>Transferrins (0.2, 1 and 5 μM, 30 min) enhances the enzymatic activity of thrombin<sup>[2]</sup>.</p> <p>?Transferrins (0.2, 1 and 5 μM, 30 min) enhances the ability of thrombin and FXIIa to hydrolyze their natural substrates (fibrinogen)<sup>[2]</sup>.</p> <p>?Transferrins can conjugate with cytochrome c, the conjugates can be uptaken by transferrin receptor overexpressing A549 cells, thereby inducing cell death<sup>[3]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

### REFERENCES

- [1]. Li H, et al. Transferrin/transferrin receptor-mediated drug delivery. Med Res Rev. 2002 May;22(3):225-50.
- [2]. Tang X, et al. Transferrin plays a central role in coagulation balance by interacting with clotting factors. Cell Res. 2020 Feb;30(2):119-132.
- [3]. Saxena M, et al. Inducing cell death in vitro in cancer cells by targeted delivery of cytochrome c via a transferrin conjugate. PLoS One. 2018 Apr 12;13(4):e0195542.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA