PEG2000-DSPE

Cat. No.: Molecular Formula: Molecular Weight: Target: Pathway:	HY-112760 C ₄₅ H ₈₇ NNaO ₁₁ P 2808.74 Liposome Metabolic Enzyme/Protease	$\sim \rho_{\rm h} \sim \rho_{\rm Ac}^{\rm pl} \sum_{\rm Nuc}^{\rm no} \rho_{\rm s} \rho_{\rm s} \sim \rho_{\rm s}^{\rm pl} \rho_{\rm s}^{\rm no} \rho_{\rm s}^{\rm pl} \rho$
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)	

SOLVENT & SOLUBILITY

		Solvent Mass Concentration	1 mg	5 mg	10 mg		
	Preparing Stock Solutions	1 mM	0.3560 mL	1.7802 mL	3.5603 mL		
		5 mM					
		10 mM					
	Please refer to the so	Please refer to the solubility information to select the appropriate solvent.					
In Vivo		 Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1.25 mg/mL (0.45 mM); Clear solution 					
		2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.25 mg/mL (0.45 mM); Clear solution					

BIOLOGICAL ACTIVITY			
Description	PEG2000-DSPE can be used for the preparation of stabilized nucleic acid-lipid particllipid particles (SNALPs). SNALPs represent some of the earliest and best functional siRNA-ABC nanoparticles described ^[1] .		

REFERENCES

[1]. Miller AD. Delivery of RNAi therapeutics: work in progress. Expert Rev Med Devices. 2013;10(6):781-811.

Product Data Sheet



Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA