Proteins



Product Data Sheet

Infliximab

Cat. No.: HY-P9970 CAS No.: 170277-31-3

Target: TNF Receptor; Bacterial Pathway: Apoptosis; Anti-infection

Storage: Please store the product under the recommended conditions in the Certificate of Analysis.

BIOLOGICAL ACTIVITY

Description	Infliximab (Avakine) is a chimeric monoclonal IgG1 antibody that specifically binds to TNF- α . Infliximab prevents the interaction of TNF- α with TNF- α receptor (TNFR1 and TNFR2). Infliximab has the potential for autoimmune, chronic inflammatory diseases and diabetic neuropathy research ^{[1][2]} .
In Vitro	TNF- α -treated adipocytes shows a significant 64% decrease in insulin-stimulated glucose uptake, whereas Infliximab (10 ng/mL) reverses TNF- α actions by significantly improving glucose incorporation in 3T3L1 adipocytes. Infliximab restores phosphorylation of substrate-2 and AKT by attenuating protein-tyrosine phosphatase 1B (PTP1B) activation. Infliximab ameliorates TNF- α -induced insulin resistance in 3T3L1 adipocytes in vitro by restoring the insulin signalling pathway via PTP1B inhibition ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	A single injection of Infliximab (10 μ g/g in 100 μ l saline/dose ip) in diabetic TNF- $\alpha^{+/+}$) mice leads to suppression of the increased serum TNF- α and amelioration of the electrophysiological and biochemical deficits for at least 4 weeks. The increased TNF- α mRNA expression in diabetic dorsal root ganglion is also attenuated by Infliximab ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- NPJ Regen Med. 2023 May 2;8(1):23.
- Int Immunopharmacol. 2023 May.
- Mol Pharm. 2022 Aug 29.
- World J Stem Cells. 2022; 14(11): 798-814.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Lucia A Méndez-García, et al. Infliximab ameliorates tumor necrosis factor-alpha-induced insulin resistance by attenuating PTP1B activation in 3T3L1 adipocytes in vitro. Scand J Immunol. 2018 Nov;88(5):e12716.

2]. Isamu Yamakawa, et al. Inactiv	ation of TNF-α ameliorates diabetic	neuropathy in mice. Am J P	hysiol Endocrinol Metab. 2011 N	ov;301(5):E844-52.
C	aution: Product has not been fu	lly validated for medical	applications. For research us	se only.
			E-mail: tech@MedChemExpre	
,		Dr, Suite Q, Monmouth J		33.00111

Page 2 of 2 www.MedChemExpress.com