


CX-5461

Cat. No.:	HY-13323		
CAS No.:	1138549-36-6		
Molecular Formula:	C ₂₇ H ₂₇ N ₇ O ₂ S		
Molecular Weight:	513.61		
Target:	DNA/RNA Synthesis		
Pathway:	Cell Cycle/DNA Damage		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : 55.56 mg/mL (108.18 mM; ultrasonic and adjust pH to 2 with HCl)
 50 mM sodium phosphate (pH 3.5) : 10 mg/mL (19.47 mM; Need ultrasonic)
 DMSO : < 1 mg/mL (ultrasonic;warming;heat to 60°C) (insoluble or slightly soluble)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	1.9470 mL	9.7350 mL	19.4700 mL
5 mM	0.3894 mL	1.9470 mL	3.8940 mL
10 mM	0.1947 mL	0.9735 mL	1.9470 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

CX-5461 is a potent and oral rRNA synthesis inhibitor. It inhibits RNA polymerase I-driven transcription of rRNA with IC₅₀s of 142, 113, and 54 nM in HCT-116, A375, and MIA PaCa-2 cells, respectively^[1].

IC₅₀ & Target

IC₅₀: 54 nM (rRNA synthesis, MIA PaCa-2 cells), 113 nM (rRNA synthesis, A375 cells), 142 nM (rRNA synthesis, HCT-116 cells)^[1]

In Vitro

CX-5461 is a potent and orally bioavailable inhibitor of Pol I-mediated rRNA synthesis, with IC₅₀s of 142 nM in HCT-116, 113 nM in A375, and 54 nM in MIA PaCa-2 cells, and shows little or no effect on Pol II (IC₅₀, ≥25 μM). CX-5461 has modest inhibition on DNA replication and protein translation. CX-5461 also exhibits broad antiproliferative activity against a panel of human cancer cell lines, with a mean EC₅₀ of 147 nM, but has minimal effect on viability of nontransformed human cells, with EC₅₀ values of appr 5000 nM. EC₅₀s of CX-5461 for HCT-116, A375, and MIA PaCa-2 cell lines are 167, 58, and 74 nM, respectively. CX-5461 induces autophagy and senescence in solid tumor cancer cells, rather than apoptosis, through a p53-independent process^[1]. Eμ-Myc lymphoma cells from tumor-bearing mice are exquisitely sensitive to CX-5461 with an IC₅₀ of 27.3 nM ± 8.1 nM for Pol I transcription after 1 hr and IC₅₀ of 5.4 nM ± 2.1 nM for cell death after 16 hr. CX-5461 activates

p53 via the nucleolar stress response in E μ -MycLymphoma Cells^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

CX-5461 displays antitumor activity against human solid tumors in murine xenograft models. CX-5461 (50 mg/kg, p.o.) shows significant MIA PaCa-2 growth inhibition with TGI equal to 69% on day 31 and 79% TGI on A375 on day 32^[1]. CX-5461 (50 mg/kg, p.o.) inhibits the E μ -Myc tumor cells with 84% repression in Pol I transcription at 1 hr posttreatment in C57BL/6 mice. CX-5461 also induces a rapid reduction in tumor burden in the lymph nodes and a concomitant reduction of spleen size to within the normal range^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

Cells are plated on 96-well plates and treated the next day with dose response of CX-5461 for 96 hours. Cell viability is determined using Alamar Blue and CyQUANT assays^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Mice^[1]

Animal experiments are performed with 5- to 6-week-old female athymic (NCr nu/nu fisol) mice of Balb/c. Mice are inoculated with athymic (NCr nu/nu fisol) mice in 100 μ L of cell suspension subcutaneously in the right flank. Tumor measurements are performed by caliper analysis, and tumor volume is calculated using the formula $(l \times w^2)/2$, where w=width and l=length in mm of the tumor. established tumors (appr 110-120 mm³) are randomized into vehicle (50 mM NaH₂PO₄, pH 4.5), NSC 613327, or CX-5461 treatment groups. Tumor growth inhibition (TGI) is determined on the last day of study according to the formula: $TGI (\%) = [100 - (Vf^D - Vi^D) / (Vf^V - Vi^V) \times 100]$, where Vi^V is the initial mean tumor volume in vehicle-treated group, Vf^V is the final mean tumor volume in vehicle-treated group, Vi^D is the initial mean tumor volume in drug-treated group, and Vf^D is the final mean tumor volume in drug-treated group.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nat Cell Biol. 2022 Jul;24(7):1154-1164.
- Nat Commun. 2022 Jun 28;13(1):3706.
- Nat Commun. 2017 Sep 25;8(1):693.
- Nucleic Acids Res. 2022 May 6;50(8):4574-4600.
- Clin Cancer Res. 2017 Nov 1;23(21):6529-6540.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Drygin D et al. Targeting RNA polymerase I with an oral small molecule CX-5461 inhibits ribosomal RNA synthesis and solid tumor growth. Cancer Res. 2011 Feb 15;71(4):1418-30.

[2]. Bywater MJ, et al. Inhibition of RNA Polymerase I as a Therapeutic Strategy to Promote Cancer-Specific Activation of p53. Cancer Cell. 2012 Jul 10;22(1):51-65.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA