

FGF basic/bFGF Protein, Human (154a.a)

Cat. No.:	HY-P7331
Synonyms:	rHubFGF, 154a.a.; bFGF; FGF-2; HBGF-2; FGFB
Species:	Human
Source:	E. coli
Accession:	P09038-4 (A135-S288)
Gene ID:	2247
Molecular Weight:	Approximately 17.1 kDa

PROPERTIES

AA Sequence	A A G S I T T L P A L P E D G G S G A F P P G H F K D P K R L Y C K N G G F F L R I H P D G R V D G V R E K S D P H I K L Q L Q A E E R G V V S I K G V C A N R Y L A M K E D G R L L A S K C V T D E C F F F E R L E S N N Y N T Y R S R K Y T S W Y V A L K R T G Q Y K L G S K T G P G Q K A I L F L P M S A K S
Biological Activity	1. The ED ₅₀ is <0.25 ng/mL as measured by 3T3 cells, corresponding to a specific activity of >4 × 10 ⁶ units/mg. 2. Measured in a cell proliferation assay using NIH3T3 cells. The ED ₅₀ for this effect is 0.1667 ng/mL, corresponding to a specific activity is 5.99×10 ⁶ units/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized after extensive dialysis against PBS, pH 7.4.
Endotoxin Level	<0.2 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	FGF-2/bFGF is a member of the fibroblast family and has a high affinity for heparin. FGF-2 plays an important role in tendon to bone healing, cartilage repair, bone repair, and nerve regeneration. FGF-2 specifically binds to tyrosine kinase receptors and activates the FGF/FGFR signaling pathway. Subsequently, FGF-2 influences cell proliferation, differentiation and apoptosis, as well as immune regulation by transducing other classical pathways. For example, FGF-2 regulates the JAK-
------------	--

STAT signaling pathway to regulate cartilage metabolism. FGF-2 also acts as a mitotic promoter to accelerate cell proliferation. Therefore, (1) FGF-2 is an important growth factor in the healing process of ligament/tendon injury. In vitro experiments, low-dose FGF-2 can stimulate the proliferation and differentiation of bone marrow mesenchymal stem cells, and up-regulate the mRNA expression of type I/III collagen and fibronectin. However, high doses of FGF-2 did not stimulate extracellular matrix (ECM) protein proliferation and gene expression. (2) FGF-2 is also an endogenous and intrinsic growth factor in cartilage repair. FGF-2 binds to heparan sulfate proteoglycan and is stored in the ECM of articular cartilage. When cartilage is damaged or degenerated, ECM rapidly releases FGF-2 and activates ERK signaling pathways to promote cartilage regeneration. FGF-2 exhibits a biphasic effect in combination with its specific receptor. FGF-2 combined with FGFR3 promoted the repair of articular cartilage. FGF-2 combined with FGFR1 promoted the degeneration of articular cartilage^[1]. FGF-2 is expressed in granulosa cells and colliculus cells, as well as hepatocellular cancer cells, but not in non-cancerous liver tissues. This reveals the role of FGF-2 in brain tumors, particularly glioblastoma. According to studies, FGF-2 is a known carcinogenic factor in GBM. FGF-2 increases the self-renewal of glioblastoma stem cells and contributes to the growth and vascularization of glioma^[2]. FGF-2 protein is highly conserved in some species, and the similarity rate of human FGF-2 protein sequence to rat, mouse, and bovine was 97.4%, 95.45%, and 98.71%, respectively.

REFERENCES

- [1]. Zhang J, et al. FGF2: a key regulator augmenting tendon-to-bone healing and cartilage repair. *Regen Med.* 2020 Sep;15(9):2129-2142.
- [2]. Jimenez-Pascual A, et al. FGF2: a novel druggable target for glioblastoma? *Expert Opin Ther Targets.* 2020 Apr;24(4):311-318.
- [3]. Hankemeier S, et al. Modulation of proliferation and differentiation of human bone marrow stromal cells by fibroblast growth factor 2: potential implications for tissue engineering of tendons and ligaments. *Tissue Eng.* 2005 Jan-Feb;11(1-2):41-9.
- [4]. Westermann R, et al. Basic fibroblast growth factor (bFGF), a multifunctional growth factor for neuroectodermal cells. *J Cell Sci Suppl.* 1990;13:97-117.
- [5]. Rusnati M, et al. Interaction of angiogenic basic fibroblast growth factor with endothelial cell heparan sulfate proteoglycans. Biological implications in neovascularization. *Int J Clin Lab Res.* 1996;26(1):15-23.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA