


CC-885

Cat. No.:	HY-101488		
CAS No.:	1010100-07-8		
Molecular Formula:	C ₂₂ H ₂₁ ClN ₄ O ₄		
Molecular Weight:	440.88		
Target:	Ligands for E3 Ligase; Molecular Glues		
Pathway:	PROTAC		
Storage:	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 67.5 mg/mL (153.10 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions			1 mg	5 mg
		1 mM		2.2682 mL	11.3410 mL
		5 mM		0.4536 mL	2.2682 mL
10 mM			0.2268 mL	1.1341 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.25 mg/mL (5.10 mM); Suspended solution; Need ultrasonic				
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.25 mg/mL (5.10 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	CC-885 is a cereblon (CRBN) modulator with potent anti-tumour activity.
IC ₅₀ & Target	CRBN ^[1] .
In Vitro	Acute myeloblastic leukemia (AML) cell lines, human liver epithelial cell line (THLE-2) and human peripheral blood mononuclear cells (PBMC) are treated with varying concentrations of CC-885, with IC ₅₀ s of 10 ⁻⁶ -1 μM. The effect of CC-885 on cell proliferation in AML cell lines, THLE-2 and human PBMC is more powerful than Lenalidomide and Pomalidomide with IC ₅₀ s > 10 μM. To address whether the cereblon-dependent degradation of GSPT1 is responsible for the cytotoxic effects of CC-885, a GSPT1 mutant that retains its normal function, but loses CC-885-dependent cereblon binding, is used to distinguish the role of GSPT1 from that of other substrates. CC-885 is tested in 293T HEK cells stably expressing the CC-885-sensitive or -resistant GSPT1 variants. Overexpression of a resistant variant GSPT1Δ(1-138)/(G575N) completely abrogate

the CC-885-induced anti-proliferation, whereas overexpression of a CC-885-sensitive variant GSPT1Δ(1-138) only confer partial protection. Similar results are obtained in AML cell lines^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

Human cancer cell lines cultured in the growth medium are seeded into black 384-well plates containing DMSO or test compounds such as CC-885 (10×10^{-6} -1 μ M). The seeding density for each cell line is optimized to allow the cell growth in the linear range during a 3-day culture period. To test the compound effect on cell proliferation in acute myeloid leukaemia (AML) cell lines, 5,000 to 10,000 cells per well in 200 μ l complete culture media are seeded into black 96-well plates containing DMSO or test compounds such as CC-885. After 48 or 72 h, cell proliferation is assessed using the CellTiter-Glo (CTG) Luminescent Cell Viability Assay^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Clin Invest. 2022 Jun 28;e153514.
- Nat Commun. 2022 Sep 10;13(1):5324.
- Cell Chem Biol. 2020 Jul 16;27(7):866-876.e8.
- Acta Pharmacol Sin. 2020 Sep;41(9):1246-1254.
- Bioorg Chem. 20 November 2021, 105505.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Mary E. Matyskiela, et al. A novel cereblon modulator recruits GSPT1 to the CRL4CRBN ubiquitin ligase. Nature. 2016 Jul 14;535(7611):252-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA